

Direct qqq Force In High Momentum Limit of QCD For Proton Physics

A.N.Mitra *

244 Tagore Park, Delhi-110009, India

Abstract

An explicit construction of the proton wave function is outlined in the high momentum limit of QCD dominated by a direct qqq force, one generated by hooking the ends of a ggg vertex to 3 distinct $\bar{q}gq$ vertices, thus making up a Y -shaped diagram (see fig.1). The high degree of S_3 symmetry thus involved ensures that the qqq wave function is a mixture of $56, 0^+$ and $20, 1^+$ components, rather than the traditional $56, 0^+$ and $70, 0^+$ type. Some results of this paradigm shift are offered.

1. Introduction

This paper (in honor of Pauchy Hwang's 60th birthday) seeks to exploit the *Symmetry theme* of this Conference for an explicit construction of the proton wave function in the high momentum limit of QCD. The theoretical framework is provided by a recent paper [1] in the background of the proton's 'physics'[2], wherein the concept of a direct 3-body force is introduced at the quark-gluon level, as a folding of a ggg vertex with 3 distinct $\bar{q}gq$ vertices, making up a Y -shaped diagram (see fig 1). As shown in [1], in the high momentum limit of QCD, where the confining qq force may be neglected, this direct qqq force *dominates* over the pairwise qq forces, thus offering a new basis for exploring the proton's wave function, one in which a full-fledged S_3 symmetry in several simultaneous d.o.f.'s plays a central role [1].

In this short presentation, we have just enough space (6 pp !) for recapitulating in barest outline the basic dynamics of ref [1], and indicate its extension for an practical construction

*Email: (1)ganmitra@nde.vsnl.net.in; (2)anmitra@physics.du.ac.in




Figure 1: (a) Pictorial view of 2- & 3- body interactions; (b) ‘Mercedes-Benz’ diagram for qqq-force

of the proton’s wave function in the high momentum limit of QCD (this part is new). This construction facilitates the effects of $SU(6)$ mixing within the dynamical framework ref [1], so as to allow for some concrete results on the proton’s ‘physics’. The dynamics is provided by a *covariant* Salpeter-like equation [3] governed by what is termed in the literature as the Markov-Yukawa Transversality Principle (MYTP for short)[4, 5] which specifies that all qq forces are *transverse* to the direction of the total hadron 4-momentum P_μ —a gauge principle in disguise [6] ! The Salpeter equation [3] has a remarkable property of a 3D-4D interlinkage [7] which can be adapted to Dirac’s LF Dynamics [8, 9] to take advantage of its bigger stability group. Using the notation, phase and normalization of ref.[1], the fully antisymmetric ‘spin’-part of the qqq force in 3D notation reads

$$V_{qqq} = \frac{-g_s^4 [SSS]}{3 k_1^2 k_2^2 k_3^2}$$

$$SSS = 2i\Sigma \cdot (\eta - \eta') \times (\xi - \xi') [k_1^2 + k_2^2 + k_3^2] / \sqrt{3} - 4[(\eta - \eta') \times (\xi - \xi')]^2 \quad (1)$$

Here Σ is twice the total hadron spin operator. The procedure is now to insert this term within an interlinked 3D-4D BSE framework wherein the complete 4D wave function Ψ satisfies a covariant Salpeter Eq a la MYTP on the light front (LF). The 4D Ψ function – the repository of all types of transition amplitudes – whose spin dependence is fully described in terms of standard Dirac matrices, can be related through a sequence of transformations to a 3D scalar function ϕ which (on reduction of the full 4D BSE for Ψ) satisfies a 3D Schroedinger-like (albeit LF covariant) equation. Next, a *reconstruction* of Ψ in terms of ϕ is achieved by Green’s function techniques, so that the 4D spin structure of Ψ is recovered. The qqq paper [1] stopped at this stage with a discussion of the general analytic structure of the 3D wave function ϕ . In this paper we seek to bridge the gap for practical applications of this formalism by interpolating between

Ψ and (ϕ) of [1], a spin-dependent 3D matrix function ψ (see below for its precise definition), which incorporates $SU(6)$ mixing effects to the extent allowed by the dynamics of V_{qqq} , Eq (1.1). The logic is similar to one employed by the Orsay group more than 3 decades ago [10], using a spin (χ)-cum-isospin (ϕ) representation for $SU(6)$ states [11], but now the dynamics of V_{qqq} automatically determines this mixing. In Sect 2, the structure of the 4D Ψ function is outlined through a sequence of Steps A,B,C of ref [1], using the 3D spin-dependent matrix function ψ as an interpolation between Ψ and ϕ . Sect 3 has just enough space to list a few results to illustrate the applications.

2. Structure of Full 4D Ψ

We collect some essential material from [1]. The LF momenta in 3D form are :

$$p_{iz}; p_{i0} = \frac{Mp_{i+}}{P_+}; \frac{Mp_{i-}}{2P_-}$$

$$\hat{p}_i \equiv \{p_{i\perp}, p_{iz}\} \quad (2.1)$$

$$\sqrt{2}\xi = p_3 - p_2; \quad \sqrt{6}\eta = -2p_3 + p_1 + p_2; \quad (2.2)$$

There are three main steps for the Ψ - ϕ interconnection

Step A: Define an auxiliary 4D scalar function Φ [1] :

$$\Psi = \Pi_{123} S_{Fi}^{-1}(-p_i) \Phi(p_i p_2 p_3) W(P) \quad (2.3)$$

where [12], [1]

$$W(P) = [\chi' \phi' + \chi'' \phi''] / \sqrt{2} \quad (2.4)$$

$$|\chi' >; |\chi'' > = \left[\frac{M - i\gamma \cdot P}{2M} [i\gamma_5; i\hat{\gamma}_\mu / \sqrt{3}] C / \sqrt{2} \right]$$

$$\otimes [[1; \gamma_5 \hat{\gamma}_\mu] u(P)] \quad (2.5)$$

Step B: Set up the Master Eq for Φ with Gordon reduction.

Step C: Make a reduction of the Master Eq for 4D Φ to one for 3D ϕ ; then *reconstruct* Φ in terms of ϕ , via Green's fn method [13] adapted to the LF formalism [1]

The final result for reconstructed 4D spinor Ψ in terms 3D scalar ϕ is

$$\Psi(\xi, \eta) = \Pi_{123} S_F(p_i) D_{123} W(P) \sum_{123} \frac{\phi(\hat{\xi}, \hat{\eta})}{(2\pi i)^2} \quad (2.6)$$

where

$$\frac{1}{D_{123}} = \int \frac{P_+^2 dq_{12-} dp_{3-}}{4M^2(2i\pi)^2 \Delta_1 \Delta_2 \Delta_3} \quad (2.7)$$

and the 3D wave fn ϕ satisfies a 6D Differential equation in coordinate space with S_3 symmetric variables

$$\sqrt{2}s_3 = r_1 - r_2; \quad \sqrt{6}t_3 = -2r_3 + r_1 + r_2 \quad (2.8)$$

Now to the specific contribution of the present paper, viz., an interpolating function ψ between Ψ and ϕ , so as to incorporate the $SU(6)$ effects. This is achieved by introducing a new 3D spin-dependent scalar matrix function ψ , via the representation :

$$\psi = \psi_s + i \frac{\sqrt{2}\Sigma \cdot \eta \times \xi}{\xi^2 + \eta^2} \psi_a \quad (2.9)$$

where ξ, η are given by Eq (2.2), and the norm of the second term is for later convenience. In the notation of ref [10], the simplest interpretation of the two scalar functions ψ_s and ψ_a is that they stand for the symmetric $(56; 0^+)$ and antisymmetric $20; 1^+$ states respectively, each with $J = 1/2$, instead of a 56-70 mixture [10]. A second difference from ref [10] is that these two functions are now *dynamically* linked by two coupled equations (c.f., the single equation for ϕ , Eq (5.18) in ref [1]). With a little approximation of angular averaging over certain terms, the ψ_s and ψ_a equations get almost decoupled and satisfy two similar equations represented symbolically as

$$D_{123}[\psi_s; \psi_a] = \int V_s[4(\eta \times \xi)^2 - 4X \pm 2X](\psi_s; \psi_a); \quad (2.10)$$

where D_{123} is given by (2.7), V_s the strength of the fully symmetric part of the qqq force, and

$$X = \rho(\eta \times \xi \cdot \hat{P})/\sqrt{3}; \quad \rho = \xi^2 + \eta^2 \quad (2.11)$$

If now as a first approximation, the (smaller) term $\pm 2X$ in Eq.(2.10) is dropped, the *ratio* of the ψ_s and ψ_0 components becomes almost independent of the dynamics, except for an overall constant ratio, while the dynamics is almost entirely contained in a common function ϕ satisfying an equation of the form (2.10). Thus

$$\psi_s = \cos \beta \phi; \quad \psi_a = \sin \beta \phi \quad (2.12)$$

where the phase factor β , plays the role of the mixing angle ϕ of ref [10], and the scalar function ϕ may now be identified with the quantity ϕ of ref.[1].




Figure 2: Schematic baryon spin diagram, with internal quark momenta p_1, p_2, p_3 ; basic spin operator $i\gamma_\mu\gamma_5$ is inserted in line p_1 .

3. Results and Discussion

Fig.2 is a generic diagram for different types of matrix elements with appropriate insertions for A . Thus the spin matrix elements g_i of ref.[2] are obtained with $A = i\gamma_\mu\gamma_5\lambda_i$ where the λ 's are the Gellmann matrixes as given in [14]. The 56-20 mixing affects g_3 but not g_8 and g_0 . The results in lowest order are

$$g_3 = (5/3) \cos^2 \beta - \sin^2 \beta; \quad g_8 = g_0 = 1/\sqrt{3} \quad (2.13)$$

Note that only g_3 depends on β , and agrees with the observed value 1.248 for $\beta \approx 24^\circ$. The other two quantities (at ≈ 0.58) agree with [2]. Of the latter, only g_8 is affected by the 2-gluon anomaly, but not g_8 .

The result for the fractional correction to $g_A^{(0)}$ due to the 2-gluon anomaly will be given in a more detailed communication on the lines of a recent preliminary analysis [15].

References

- [1] A. N.Mitra, Ann Phys(NY) **323**, 845-865 (2007).
- [2] S.D. Bass Rev. Mod. Phys. **77**, 1257-1302 (2005).
- [3] E.E. Salpeter, Phys.Rev. **87** (1952) 328.
- [4] M.A. Markov, Sov. J. Phys. **3** (1940) 452.
- [5] H. Yukawa, Phys.Rev. **77**(1950) 219.

- [6] J. Lukierski and M. Oziwicz, Phys. Lett.**B69**, 339 (1977).
- [7] A.N. Mitra, Nucl. Phys. **A684**, 313c (2000)
- [8] P.A.M. Dirac, Rev. Mod. Phys. **21** (1949) 392.
- [9] S. Weinberg, Phys. Rev. **150** (1966) 1313
- [10] A.Le Yaouanc et al, Phys.Rev.**D12** (1975) 2137.
- [11] A.N. Mitra and M.H. Ross, Phys. Rev.**158**, 1630 (1967)
- [12] R. Blankenbecler et al, Nucl.Phys.**12**, 629 (1959)
- [13] A.N. Mitra Int J Mod Phys.**A14**, 4781 (1999).
- [14] D.B. Lichtenberg, Unitary Symmetry and Elementary Particles, Academic Press N.Y. (1978).
- [15] A.N. Mitra, hep-ph 0712.2896